

# THE JOINT ESSENTIAL NUMERICAL RANGE OF OPERATORS: CONVEXITY AND RELATED RESULTS

CHI-KWONG LI <sup>\*</sup>and YIU-TUNG POON<sup>†</sup>

## Abstract

Let  $W(\mathbf{A})$  and  $W_e(\mathbf{A})$  be the joint numerical range and the joint essential numerical range of an  $m$ -tuple of self-adjoint operators  $\mathbf{A} = (A_1, \dots, A_m)$  acting on an infinite dimensional Hilbert space, respectively. In this paper, it is shown that  $W_e(\mathbf{A})$  is always convex and admits many equivalent formulations. In particular, for any fixed  $i \in \{1, \dots, m\}$ ,  $W_e(\mathbf{A})$  can be obtained as the intersection of all sets of the form

$$\mathbf{cl}(W(A_1, \dots, A_{i+1}, A_i + F, A_{i+1}, \dots, A_m)),$$

where  $F = F^*$  has finite rank. Moreover, it is shown that the closure  $\mathbf{cl}(W(\mathbf{A}))$  of  $W(\mathbf{A})$  is always star-shaped with the elements in  $W_e(\mathbf{A})$  as star centers. Although  $\mathbf{cl}(W(\mathbf{A}))$  is usually not convex, an analog of the separation theorem is obtained, namely, for any element  $\mathbf{d} \notin \mathbf{cl}(W(\mathbf{A}))$ , there is a linear functional  $f$  such that  $f(\mathbf{d}) > \sup\{f(\mathbf{a}) : \mathbf{a} \in \mathbf{cl}(W(\tilde{\mathbf{A}}))\}$ , where  $\tilde{\mathbf{A}}$  is obtained from  $\mathbf{A}$  by perturbing one of the components  $A_i$  by a finite rank self-adjoint operator. Other results on  $W(\mathbf{A})$  and  $W_e(\mathbf{A})$  extending those on a single operator are obtained.

**AMS Subject Classification** 47A12, 47A13, 47A55.

**Keywords** Joint essential numerical range, self-adjoint operator, Hilbert space.

---

<sup>\*</sup>Department of Mathematics, The College of William and Mary, Williamsburg, Virginia 23185, USA (ckli@math.wm.edu). Li is an honorary professor of the University of Hong Kong. His research was partially supported by an NSF grant and the William and Mary Plumeri Award

<sup>†</sup>Department of Mathematics, Iowa State University, Ames, IA (ytpoon@iastate.edu).

# 1 Introduction

Let  $\mathcal{B}(\mathcal{H})$  denote the algebra of bounded linear operators acting on a complex Hilbert space  $\mathcal{H}$ . The *numerical range* of  $A \in \mathcal{B}(\mathcal{H})$  is defined as

$$W(A) = \{\langle A\mathbf{x}, \mathbf{x} \rangle : \mathbf{x} \in \mathcal{H}, \langle \mathbf{x}, \mathbf{x} \rangle = 1\},$$

which is useful in studying operators; see [10, 11, 22, 24] and [25, Chapter 1]. Let  $\mathcal{S}(\mathcal{H})$  denote the set of self-adjoint operators in  $\mathcal{B}(\mathcal{H})$ . Since every  $A \in \mathcal{B}(\mathcal{H})$  admits a decomposition  $A = A_1 + iA_2$  with  $A_1, A_2 \in \mathcal{S}(\mathcal{H})$ , we can identify  $W(A)$  with

$$\{(\langle A_1\mathbf{x}, \mathbf{x} \rangle, \langle A_2\mathbf{x}, \mathbf{x} \rangle) : \mathbf{x} \in \mathcal{H}, \langle \mathbf{x}, \mathbf{x} \rangle = 1\} \subseteq \mathbf{R}^2.$$

This leads to the *joint numerical range* of  $\mathbf{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ ,

$$W(\mathbf{A}) = \{(\langle A_1\mathbf{x}, \mathbf{x} \rangle, \dots, \langle A_m\mathbf{x}, \mathbf{x} \rangle) : \mathbf{x} \in \mathcal{H}, \langle \mathbf{x}, \mathbf{x} \rangle = 1\} \subseteq \mathbf{R}^m,$$

which has been studied by many researchers in order to understand the joint behavior of several operators  $A_1, \dots, A_m$ . One may see [1, 5, 12, 14, 15, 16, 19, 23, 28, 31, 33, 35] and their references for the background and many applications of the joint numerical range.

Let  $\mathcal{F}(\mathcal{H})$  and  $\mathcal{K}(\mathcal{H})$  be the sets of finite rank and compact operators in  $\mathcal{B}(\mathcal{H})$ . In the study of finite rank or compact perturbations of operators, researchers consider the *joint essential numerical range* of  $\mathbf{A} \in \mathcal{S}(\mathcal{H})^m$  defined by

$$W_e(\mathbf{A}) = \cap \{\mathbf{cl}(W(\mathbf{A} + \mathbf{K})) : \mathbf{K} = (K_1, \dots, K_m) \in \mathcal{K}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m\}.$$

Here  $\mathbf{cl}(S)$  denotes the closure of the set  $S$ . For  $m = 2$ ,  $W_e(\mathbf{A})$  can be identified with the *essential numerical range* of  $A = A_1 + iA_2 \in \mathcal{B}(\mathcal{H})$  defined by

$$W_e(A) = \cap \{\mathbf{cl}(W(A + K)) : K \in \mathcal{K}(\mathcal{H})\}.$$

One may see [2, 3, 6, 7, 13, 18, 20, 21, 26, 27, 30, 32, 36, 37] for many interesting results on  $W_e(A)$  and  $W_e(\mathbf{A})$ .

In theoretical study as well as applications, it is desirable to deal with  $\mathbf{A}$  such that  $W(\mathbf{A})$  or  $\mathbf{cl}(W(\mathbf{A}))$  is convex. For example, let  $\mathbf{A} = (A_1, \dots, A_m)$ . If  $\mathbf{cl}(W(\mathbf{A}))$  is convex, one can apply the separation theorem to show that  $\mathbf{0} \notin \mathbf{cl}(W(\mathbf{A}))$  if and only if there exist  $r > 0$  and  $\mathbf{c} = (c_1, \dots, c_m) \in \mathbf{R}^m$  such that  $(\sum_{i=1}^m c_i A_i) > rI_{\mathcal{H}}$ . Unfortunately,  $\mathbf{cl}(W(\mathbf{A}))$  is not always convex. Here are some results concerning the convexity of  $W(\mathbf{A})$  and  $\mathbf{cl}(W(\mathbf{A}))$ , and related to  $W_e(\mathbf{A})$ ; for example, see [5, 10, 11, 36, 21, 29, 31] and their references.

- (P1) [31]  $W(A_1, \dots, A_m)$  is convex if
- (a)  $\text{span}\{I, A_1, \dots, A_m\}$  has dimension at most 3, or
  - (b)  $\dim \mathcal{H} \geq 3$  and  $\text{span}\{I, A_1, \dots, A_m\}$  has dimension at most 4.
- (P2) [31] For any  $A_1, A_2, A_3 \in \mathcal{S}(\mathcal{H})$  such that  $\text{span}\{I, A_1, A_2, A_3\}$  has dimension 4, there is always an  $A_4 \in \mathcal{S}(\mathcal{H})$  for which  $W(A_1, \dots, A_4)$  is not convex.
- (P3) [31] If  $m \geq 4$  then there exists  $\mathbf{A} \in \mathcal{S}(\mathcal{H})^m$  such that  $W(\mathbf{A})$  is non-convex.
- (P4) For any positive integer  $m$  and any  $\mathbf{A} \in \mathcal{S}(\mathcal{H})^m$ ,  $W_e(\mathbf{A})$  is a compact set contained in  $W(\mathbf{A})$ . If  $\text{span}\{I, A_1, \dots, A_m\}$  has dimension at most 4, then  $W_e(\mathbf{A})$  is convex.
- (P5) [36] For  $S \subseteq \mathbf{R}^m$ , let  $\text{Ext}(S)$  be the set of all points in  $S$  that does not lie in the open line segment joining two distinct points in  $S$ . Then  $\text{Ext}(\text{cl}(W(\mathbf{A}))) \subseteq \text{Ext}(W(\mathbf{A})) \cup \text{Ext}(W_e(\mathbf{A}))$ .

We remark that (P1)-(P3) also hold if we replace  $W(\mathbf{A})$  by  $\text{cl}(W(\mathbf{A}))$ . In view of (P2) and (P3), if  $m > 3$ , then for  $\mathbf{A} \in \mathcal{S}(\mathcal{H})^m$  and  $\mathbf{K} \in \mathcal{K}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m$  the set  $\text{cl}(W(\mathbf{A} + \mathbf{K}))$  is usually non-convex. Since  $W_e(\mathbf{A})$  is the intersection of non-convex sets, one does not expect the set  $W_e(\mathbf{A})$  to be convex. This might be the reason why the convexity of  $W_e(\mathbf{A})$  is seldom discussed for  $m > 3$ . In fact, some researchers have studied different geometrical properties of  $W_e(\mathbf{A})$  under the assumption that  $W_e(\mathbf{A})$  is convex, and some researchers studied  $W_e(\mathbf{A})$  for different classes of operators without discussing their convexity; for example, see [6, 26, 27, 30, 32].

In this paper, we prove the rather unexpected result that  $W_e(\mathbf{A})$  is *always convex*. Moreover, it is shown that the closure  $\text{cl}(W(\mathbf{A}))$  of  $W(\mathbf{A})$  is always star-shaped with the elements in  $W_e(\mathbf{A})$  as star centers. Many results relating  $W_e(\mathbf{A})$  and  $W(\mathbf{A})$  are also obtained. Our paper is organized as follows.

In Section 2, we extend the results in [21] to establish several equivalent formulations of the essential joint numerical range for  $\mathbf{A} \in \mathcal{S}(\mathcal{H})^m$ . One key obstacle for such an extension is the fact that  $W(\mathbf{A})$  may not be convex. To get around this problem, we show that  $\text{cl}(W(\mathbf{A}))$  is star-shaped. The star-shapedness of  $\text{cl}(W(\mathbf{A}))$  and the equivalent formulations of  $W_e(\mathbf{A})$  in Section 2 lead to our main result that  $W_e(\mathbf{A})$  is convex and its elements are

star centers of the set  $\mathbf{cl}(W(\mathbf{A}))$ , which is presented in Section 3. With the convexity theorem, we obtain additional descriptions of  $W_e(\mathbf{A})$  in Section 4 in terms of the perturbations of one of the components of  $\mathbf{A}$ , and also in terms of linear combinations of the components of  $\mathbf{A}$ . For example, we show that  $W_e(A_1, \dots, A_m)$  is equal to the sets

$$\cap\{\mathbf{cl}(W(A_1, \dots, A_{i-1}, A_i + F, A_{i+1}, \dots, A_m)) : F \in \mathcal{F}(\mathcal{H}) \cap \mathcal{S}(\mathcal{H})\}$$

and

$$\left\{ (a_1, \dots, a_m) : \sum_{j=1}^m c_j a_j \in W_e \left( \sum_{j=1}^m c_j A_j \right) \text{ for all } (c_1, \dots, c_m) \in \Omega \right\},$$

where  $\Omega = \{(c_1, \dots, c_m) \in \mathbf{R}^m : \sum_{j=1}^m c_j^2 = 1\}$ . Also, we obtain an analog of the separation theorem for the not necessarily convex set  $\mathbf{cl}(W(\mathbf{A}))$ , namely, for any element  $\mathbf{d} \notin \mathbf{cl}(W(\mathbf{A}))$ , there is a linear functional  $f$  such that  $f(\mathbf{d}) > \sup\{f(\mathbf{a}) : \mathbf{a} \in \mathbf{cl}(W(\tilde{\mathbf{A}}))\}$ , where  $\tilde{\mathbf{A}}$  is obtained from  $\mathbf{A}$  by perturbing one of the components  $A_j$  by a finite rank self-adjoint operator. In Section 5, we present additional results on  $W(\mathbf{A})$  and  $W_e(\mathbf{A})$ . For instance,  $W_e(\mathbf{A}) = \mathbf{cl}(W(\mathbf{A}))$  if and only if the extreme points of  $W(\mathbf{A})$  are contained in  $W_e(\mathbf{A})$ ; the convex hull of  $\mathbf{cl}(W(\mathbf{A}))$  can always be realized the the joint essential numerical range of  $(\tilde{A}_1, \dots, \tilde{A}_m)$  for linear operators  $\tilde{A}_1, \dots, \tilde{A}_m$  acting on a separable Hilbert space.

In our discussion, we always assume that  $\mathcal{H}$  is infinite-dimensional. For any vector  $\mathbf{x} \in \mathcal{H}$  and  $\mathbf{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ , we will use the following notation

$$\langle \mathbf{A}\mathbf{x}, \mathbf{x} \rangle = (\langle A_1\mathbf{x}, \mathbf{x} \rangle, \dots, \langle A_m\mathbf{x}, \mathbf{x} \rangle).$$

Furthermore,  $\mathbf{R}^m$  will be used to denote the inner product space of  $1 \times m$  real vectors with the usual inner product  $\langle \mathbf{x}, \mathbf{y} \rangle$ .

## 2 Equivalent formulations of $W_e(\mathbf{A})$

Following [21, Theorem 5.1] and its corollary on a single operator  $A \in \mathcal{B}(\mathcal{H})$ , we obtain several equivalent formulations of  $W_e(\mathbf{A})$ .

**Theorem 2.1** *Let  $\mathbf{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ . The following conditions are equivalent for a real vector  $\mathbf{a} = (a_1, \dots, a_m)$ .*

- (1)  $\mathbf{a} \in W_e(\mathbf{A}) = \cap\{\mathbf{cl}(W(\mathbf{A} + \mathbf{K})) : \mathbf{K} \in \mathcal{K}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m\}$ .
- (2)  $\mathbf{a} \in \cap\{\mathbf{cl}(W(\mathbf{A} + \mathbf{F})) : \mathbf{F} \in \mathcal{F}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m\}$ .

(3) There is an orthonormal sequence of vectors  $\{\mathbf{x}_n\}_{n=1}^\infty \in \mathcal{H}$  such that

$$\lim_{n \rightarrow \infty} \langle \mathbf{A}\mathbf{x}_n, \mathbf{x}_n \rangle = \mathbf{a}.$$

(4) There is a sequence of unit vectors  $\{\mathbf{x}_n\}_{n=1}^\infty \in \mathcal{H}$  converging weakly to  $\mathbf{0}$  in  $\mathcal{H}$  such that

$$\lim_{n \rightarrow \infty} \langle \mathbf{A}\mathbf{x}_n, \mathbf{x}_n \rangle = \mathbf{a}.$$

(5) There is an infinite-dimensional projection  $P \in \mathcal{S}(\mathcal{H})$  such that

$$P(A_j - a_j I)P \in \mathcal{K}(\mathcal{H}) \quad \text{for } j = 1, \dots, k.$$

Most of the argument in [21] can be applied here except for one crucial step, where the convexity of  $W(\mathbf{A})$  for  $m = 2$  is needed. Since  $W(\mathbf{A})$  may not be convex for  $m > 3$ , we need the following auxiliary result to overcome the obstacle. As a byproduct, it shows that  $\mathbf{cl}(W(\mathbf{A}))$  is star-shaped.

**Theorem 2.2** *Let  $\mathbf{A}$  satisfy the hypothesis of Theorem 2.1, and let  $W_3(\mathbf{A})$  be the set of real vectors  $\mathbf{a}$  satisfying condition (3) of Theorem 2.1. Then  $W_3(\mathbf{A})$  is non-empty and closed. Moreover, each element  $\mathbf{a} \in W_3(\mathbf{A})$  is a star center of  $\mathbf{cl}(W(\mathbf{A}))$ , i.e., for any  $\mathbf{b} \in \mathbf{cl}(W(\mathbf{A}))$  we have  $(1-t)\mathbf{a} + t\mathbf{b} \in \mathbf{cl}(W(\mathbf{A}))$  for all  $0 \leq t \leq 1$ .*

*Proof.* To prove that  $W_3(\mathbf{A})$  is non-empty, let  $\{\mathbf{x}_n\}_{n=1}^\infty$  be an orthonormal sequence of vectors in  $\mathcal{H}$ . Then the sequence  $\{\langle \mathbf{A}\mathbf{x}_n, \mathbf{x}_n \rangle\}_{n=1}^\infty$  is bounded. By choosing a subsequence, if necessary, we can assume that  $\langle \mathbf{A}\mathbf{x}_n, \mathbf{x}_n \rangle$  converges. Hence,  $W_3(\mathbf{A})$  is non-empty.

Next, we show that  $W_3(\mathbf{A})$  is closed. Suppose  $\mathbf{a} \in \mathbf{cl}(W_3(\mathbf{A}))$ . Then for each  $n \geq 1$ , there exists an orthonormal sequence  $\{\mathbf{x}_k^n\}_{k=1}^\infty$  such that  $\lim_{k \rightarrow \infty} \langle \mathbf{A}\mathbf{x}_k^n, \mathbf{x}_k^n \rangle = \mathbf{a}^n \in \mathbf{R}^m$  and  $\lim_{n \rightarrow \infty} \mathbf{a}^n = \mathbf{a}$ . Let  $\delta_n = 1/(4n^2)$ . By going to subsequences, if necessary, we may assume that  $\|\langle \mathbf{A}\mathbf{x}_k^n, \mathbf{x}_k^n \rangle - \mathbf{a}\| < \delta_n$  for all  $n, k$ . We may also assume that  $\|A_1\|^2 + \dots + \|A_m\|^2 \leq 1$ . Then  $\|\langle \mathbf{A}\mathbf{x}, \mathbf{y} \rangle\| \leq \|\mathbf{x}\|\|\mathbf{y}\|$  for all  $\mathbf{x}, \mathbf{y} \in \mathcal{H}$ .

Choose  $\mathbf{x}_1 = \mathbf{x}_1^1$ . Then we have  $\|\langle \mathbf{A}\mathbf{x}_1, \mathbf{x}_1 \rangle - \mathbf{a}\| < 1$ . Suppose we have chosen  $\{\mathbf{x}_1, \dots, \mathbf{x}_n\}$  orthonormal, with  $\|\langle \mathbf{A}\mathbf{x}_k, \mathbf{x}_k \rangle - \mathbf{a}\| < 1/k$  for  $1 \leq k \leq n$ . Then choose  $N$  such that for all  $1 \leq k \leq n$ , we have

$$|\langle \mathbf{x}_k, \mathbf{x}_N^{n+1} \rangle|, \|\langle \mathbf{A}\mathbf{x}_k, \mathbf{x}_N^{n+1} \rangle\| < \delta_{n+1}.$$

Let  $\mathbf{y} = \mathbf{x}_N^{n+1} - \sum_{k=1}^n \langle \mathbf{x}_N^{n+1}, \mathbf{x}_k \rangle \mathbf{x}_k$ . Then

$$\|\mathbf{y} - \mathbf{x}_N^{n+1}\| \leq n\delta_{n+1} \Rightarrow 1 - n\delta_{n+1} \leq \|\mathbf{y}\| \leq 1 + n\delta_{n+1}.$$

Therefore,

$$\begin{aligned}
& \| \langle \mathbf{A}\mathbf{y}, \mathbf{y} \rangle - \mathbf{a} \| \\
& \leq \| \langle \mathbf{A}(\mathbf{y} - \mathbf{x}_N^{n+1}), \mathbf{y} \rangle \| + \| \langle \mathbf{A}\mathbf{x}_N^{n+1}, \mathbf{y} - \mathbf{x}_N^{n+1} \rangle \| + \| \langle \mathbf{A}\mathbf{x}_N^{n+1}, \mathbf{x}_N^{n+1} \rangle - \mathbf{a} \| \\
& \leq \| \mathbf{y} - \mathbf{x}_N^{n+1} \| (\| \mathbf{y} \| + \| \mathbf{x}_N^{n+1} \|) + \delta_{n+1} \\
& \leq (2n + 2)\delta_{n+1}.
\end{aligned}$$

Let  $\mathbf{x}_{n+1} = \mathbf{y}/\| \mathbf{y} \|$ . Then

$$\| \mathbf{x}_{n+1} - \mathbf{y} \| = |1 - \| \mathbf{y} \| | \leq n\delta_{n+1}.$$

Hence,  $\{ \mathbf{x}_1, \dots, \mathbf{x}_n, \mathbf{x}_{n+1} \}$  is an orthonormal set and

$$\begin{aligned}
\| \langle \mathbf{A}\mathbf{x}_{n+1}, \mathbf{x}_{n+1} \rangle - \mathbf{a} \| & \leq \| \mathbf{y} - \mathbf{x}_{n+1} \| (\| \mathbf{y} \| + \| \mathbf{x}_{n+1} \|) + (2n + 2)\delta_{n+1} \\
& \leq (4n + 3)\delta_{n+1} < 1/(n + 1).
\end{aligned}$$

To prove the last assertion, let  $\mathbf{a} \in W_3(\mathbf{A})$  and  $\mathbf{b} \in \mathbf{cl}(W(\mathbf{A}))$ . Suppose  $\{ \mathbf{x}_n \}$  is an orthonormal sequence in  $\mathcal{H}$  such that  $\langle \mathbf{A}\mathbf{x}_n, \mathbf{x}_n \rangle \rightarrow \mathbf{a}$ . For  $0 \leq t \leq 1$ , we are going to show that  $(1 - t)\mathbf{a} + t\mathbf{b} \in \mathbf{cl}(W(\mathbf{A}))$ . Given  $\varepsilon > 0$ , let  $\mathbf{y}$  be a unit vector in  $\mathcal{H}$  such that  $\| \langle \mathbf{A}\mathbf{y}, \mathbf{y} \rangle - \mathbf{b} \| < \varepsilon$ . Choose  $n$  such that  $\| \langle \mathbf{A}\mathbf{x}_n, \mathbf{x}_n \rangle - \mathbf{a} \| < \varepsilon$  and  $\| \langle \mathbf{A}\mathbf{y}, \mathbf{x}_n \rangle \| < \varepsilon$ . Choose  $\theta \in \mathbf{R}$  such that  $\langle e^{i\theta}\mathbf{y}, \mathbf{x}_n \rangle$  is imaginary. Let  $\mathbf{z} = \sqrt{t}e^{i\theta}\mathbf{y} + \sqrt{1-t}\mathbf{x}_n$ . Then we have

$$\langle \mathbf{z}, \mathbf{z} \rangle = t\langle \mathbf{y}, \mathbf{y} \rangle + (1 - t)\langle \mathbf{x}_n, \mathbf{x}_n \rangle + 2\sqrt{t}\sqrt{1-t}(\langle e^{i\theta}\mathbf{y}, \mathbf{x}_n \rangle + \langle \mathbf{x}_n, e^{i\theta}\mathbf{y} \rangle) = 1$$

and

$$\begin{aligned}
& \| \langle \mathbf{A}\mathbf{z}, \mathbf{z} \rangle - ((1 - t)\mathbf{a} + t\mathbf{b}) \| \\
& \leq (1 - t)\| \langle \mathbf{A}\mathbf{x}_n, \mathbf{x}_n \rangle - \mathbf{a} \| + t\| \langle \mathbf{A}\mathbf{y}, \mathbf{y} \rangle - \mathbf{b} \| \\
& \quad + \sqrt{t}\sqrt{1-t}\| \langle e^{i\theta}\mathbf{A}\mathbf{y}, \mathbf{x}_n \rangle + \langle \mathbf{A}\mathbf{x}_n, e^{i\theta}\mathbf{y} \rangle \| \\
& \leq 2\varepsilon.
\end{aligned}$$

Therefore,  $(1 - t)\mathbf{a} + t\mathbf{b} \in \mathbf{cl}(W(\mathbf{A}))$ . □

The referee indicated that  $W_3(\mathbf{A})$  is clearly closed, and a short proof is possible. We include the detailed proof for the sake of completeness and easy reference.

*Proof of Theorem 2.1.* For  $j = 2, 3, 4, 5$ , let  $W_j(\mathbf{A})$  be the set of  $\mathbf{a}$  satisfying condition (j). Clearly, we have

$$W_5(\mathbf{A}) \subseteq W_3(\mathbf{A}) \subseteq W_4(\mathbf{A}) \subseteq W_e(\mathbf{A}) \subseteq W_2(\mathbf{A}).$$

Suppose  $\mathbf{a} \in W_2(\mathbf{A})$ . We are going to show that  $\mathbf{a} \in W_5(\mathbf{A})$ . Without loss of generality, we may assume  $\mathbf{a} = \mathbf{0}$ .

Since  $\mathbf{0} \in W_2(\mathbf{A}) \subseteq \mathbf{cl}(W(\mathbf{A}))$ , there exists a unit vector  $\mathbf{x}_1 \in \mathcal{H}$  such that  $\|\langle \mathbf{A}\mathbf{x}_1, \mathbf{x}_1 \rangle\| < 1/2$ . Suppose we have an orthonormal set of vectors  $\{\mathbf{x}_1, \dots, \mathbf{x}_n\}$  such that  $\|\langle \mathbf{A}\mathbf{x}_n, \mathbf{x}_n \rangle\| < 1/2^n$ . Let  $Q$  be the orthogonal projection of  $\mathcal{H}$  onto the subspace  $S$  spanned by  $\mathbf{x}_1, \dots, \mathbf{x}_n$  and

$$\mathbf{B} = ((I - Q)A_1(I - Q)|_{S^\perp}, \dots, (I - Q)A_m(I - Q)|_{S^\perp}).$$

Let  $\mathbf{b} = (b_1, \dots, b_m) \in W_3(\mathbf{B})$  and  $\mathbf{b}I_S = (b_1I_S, \dots, b_mI_S)$ . Then for  $\bar{Q} = I - Q$ , we have

$$\mathbf{b}I_S \oplus \mathbf{B} = (b_1Q + \bar{Q}A_1\bar{Q}, \dots, b_mQ + \bar{Q}A_m\bar{Q}) = \mathbf{A} + \mathbf{F}$$

for some  $\mathbf{F} \in \mathcal{F}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m$ . Therefore,  $\mathbf{0} \in \mathbf{cl}(W(\mathbf{b}I_S \oplus \mathbf{B}))$ . Hence, there exists a unit vector  $\mathbf{x} \in \mathcal{H}$  such that  $\|\langle (\mathbf{A} + \mathbf{F})\mathbf{x}, \mathbf{x} \rangle\| < 1/(2^{n+2})$ . Let  $\mathbf{x} = \mathbf{y} + \mathbf{z}$ , where  $\mathbf{y} \in S$  and  $\mathbf{z} \in S^\perp$ . Then  $\|\mathbf{y}\|^2 + \|\mathbf{z}\|^2 = \|\mathbf{x}\|^2 = 1$ . If  $\mathbf{z} = \mathbf{0}$ , then  $\langle (\mathbf{A} + \mathbf{F})\mathbf{x}, \mathbf{x} \rangle = \mathbf{b} \in W_3(\mathbf{B}) \subseteq \mathbf{cl}(W(\mathbf{B}))$ . If  $\mathbf{z} \neq \mathbf{0}$ , then by Theorem 2.2, we have

$$\langle (\mathbf{A} + \mathbf{F})\mathbf{x}, \mathbf{x} \rangle = \|\mathbf{y}\|^2\mathbf{b} + \|\mathbf{z}\|^2\langle \mathbf{B} \left( \frac{\mathbf{z}}{\|\mathbf{z}\|} \right), \left( \frac{\mathbf{z}}{\|\mathbf{z}\|} \right) \rangle \in \mathbf{cl}(W(\mathbf{B})).$$

So there exists a unit vector  $\mathbf{x}_{n+1} \in S^\perp$  such that

$$\|\langle (\mathbf{A} + \mathbf{F})\mathbf{x}, \mathbf{x} \rangle - \langle \mathbf{B}\mathbf{x}_{n+1}, \mathbf{x}_{n+1} \rangle\| < \frac{1}{2^{n+2}}$$

$$\Rightarrow \|\langle \mathbf{A}\mathbf{x}_{n+1}, \mathbf{x}_{n+1} \rangle\| = \|\langle \mathbf{B}\mathbf{x}_{n+1}, \mathbf{x}_{n+1} \rangle\| < \frac{1}{2^{n+1}},$$

because  $\langle \mathbf{F}\mathbf{x}_{n+1}, \mathbf{x}_{n+1} \rangle = \mathbf{0}$ . Inductively, we can choose an orthonormal sequence of vectors  $\{\mathbf{x}_n\}_{n=1}^\infty$  such that

$$\|\langle \mathbf{A}\mathbf{x}_n, \mathbf{x}_n \rangle\| < \frac{1}{2^n} \quad \text{for all } n \geq 1. \quad (1)$$

Let  $n_1 = 1$ . For every  $1 \leq i \leq m$ , we have

$$\sum_{n=1}^{\infty} |\langle A_i\mathbf{x}_{n_1}, \mathbf{x}_n \rangle|^2 \leq \|A_i\mathbf{x}_{n_1}\|^2 \quad \text{and} \quad \sum_{n=1}^{\infty} |\langle A_i\mathbf{x}_n, \mathbf{x}_{n_1} \rangle|^2 \leq \|A_i^*\mathbf{x}_{n_1}\|^2.$$

Hence, there exists  $n_2 > n_1$  such that

$$\sum_{n=n_2}^{\infty} |\langle A_i\mathbf{x}_{n_1}, \mathbf{x}_n \rangle|^2 < 1/2 \quad \text{and} \quad \sum_{n=n_2}^{\infty} |\langle A_i\mathbf{x}_n, \mathbf{x}_{n_1} \rangle|^2 < 1/2$$

for all  $1 \leq i \leq m$ . Repeating this procedure, we can get a strictly increasing sequence  $\{n_k\}_{k=1}^{\infty}$  of positive integers such that for all  $1 \leq i \leq m$ , we have

$$\sum_{n=n_{k+1}}^{\infty} |\langle A_i \mathbf{x}_{n_k}, \mathbf{x}_n \rangle|^2 < 1/2^k \quad \text{and} \quad \sum_{n=n_{k+1}}^{\infty} |\langle A_i \mathbf{x}_n, \mathbf{x}_{n_k} \rangle|^2 < 1/2^k. \quad (2)$$

(1) and (2) imply that

$$\sum_{k,\ell=1}^{\infty} |\langle A_i \mathbf{x}_{n_k}, \mathbf{x}_{n_\ell} \rangle|^2 < \infty. \quad (3)$$

Let  $P$  be the orthogonal projection onto the subspace spanned by  $\{\mathbf{x}_{n_k}\}_{k=1}^{\infty}$ . Then it follows from (3) that  $PA_iP$  is compact for all  $1 \leq i \leq m$ .  $\square$

### 3 Convexity and star-shapedness

**Theorem 3.1** *Let  $\mathbf{A} \in \mathcal{S}(\mathcal{H})^m$ . Then  $W_e(\mathbf{A})$  is a compact convex subset of  $\text{cl}(W(\mathbf{A}))$ . Moreover, each element in  $W_e(\mathbf{A})$  is a star center of the star-shaped set  $\text{cl}(W(\mathbf{A}))$ .*

*Proof.* Because  $W_e(\mathbf{A})$  is the intersection of compact sets, it is compact. To prove the convexity, let  $\mathbf{a}, \mathbf{b} \in W_e(\mathbf{A})$  and  $0 \leq t \leq 1$ . Then for every  $\mathbf{F} \in \mathcal{F}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m$ , we have  $\mathbf{a} \in W_e(\mathbf{A}) = W_e(\mathbf{A} + \mathbf{F})$  and  $\mathbf{b} \in W_e(\mathbf{A}) \subseteq \text{cl}(W(\mathbf{A} + \mathbf{F}))$ . So, by Theorem 2.2, we have  $t\mathbf{a} + (1-t)\mathbf{b} \in \text{cl}(W(\mathbf{A} + \mathbf{F}))$ . Hence,

$$t\mathbf{a} + (1-t)\mathbf{b} \in \cap \{\text{cl}(W(\mathbf{A} + \mathbf{F})) : \mathbf{F} \in \mathcal{F}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m\} = W_e(\mathbf{A}).$$

By Theorem 2.1 and Theorem 2.2, we have the last assertion.  $\square$

Note that  $W_e(\mathbf{A}) \cap W(\mathbf{A})$  may be empty. For example, if

$$A = \text{diag}(1, 1/2, 1/3, \dots)$$

acts on  $\ell^2$ , then  $W_e(A) = \{0\}$  and  $W(A) = (0, 1]$ . One may wonder whether a point  $\mathbf{a} \in W_e(\mathbf{A}) \cap W(\mathbf{A})$  is a star center of  $W(\mathbf{A})$ . This is not true as shown by the following example. Moreover, the example shows that for  $m \geq 4$  there exists  $\mathbf{A} \in \mathcal{S}(\mathcal{H})^m$  such that  $\text{cl}(W(\mathbf{A}))$  is convex whereas  $W(\mathbf{A})$  is not. Of course this is impossible for  $m \leq 3$  as  $W(\mathbf{A})$  is always convex.

**Example 3.2** Consider  $\mathcal{H} = \ell^2$  with canonical basis  $\{e_n : n \geq 1\}$ . Let  $\mathbf{A} = (A_1, \dots, A_4)$  with  $A_1 = \text{diag}(1, 0, 1/3, 1/4, \dots)$ ,  $A_2 = \text{diag}(1, 0) \oplus \mathbf{0}$ ,

$$A_3 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus \mathbf{0} \quad \text{and} \quad A_4 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} \oplus \mathbf{0}.$$

Then  $(1, 1, 0, 0) \in W(\mathbf{A})$  and  $(0, 0, 0, 0) \in W(\mathbf{A}) \cap W_e(\mathbf{A})$ , but  $(1/2, 1/2, 0, 0) \notin W(\mathbf{A})$ . Hence,  $W(\mathbf{A})$  is not convex. However,  $\text{cl}(W(\mathbf{A}))$  is convex.

*Proof of the claims in the example.* Note that  $(1, 1, 0, 0) = \langle \mathbf{A}e_1, e_1 \rangle \in W(\mathbf{A})$  and

$$(0, 0, 0, 0) = \langle \mathbf{A}e_2, e_2 \rangle = \lim_{n \rightarrow \infty} \langle \mathbf{A}e_n, e_n \rangle \in W(\mathbf{A}) \cap W_e(\mathbf{A}).$$

To show that  $(1/2, 1/2, 0, 0) \notin W(\mathbf{A})$ , consider a unit vector  $\mathbf{x} = \sum x_j e_j$  such that  $\sum_{n=1}^{\infty} |x_n|^2 = 1$ . If  $\langle A_1 \mathbf{x}, \mathbf{x} \rangle = \langle A_2 \mathbf{x}, \mathbf{x} \rangle = 1/2$ , then

$$|x_1|^2 + \sum_{n=3}^{\infty} |x_n|^2/n = |x_1|^2 = 1/2.$$

Thus,  $x_n = 0$  for all  $n \geq 3$  and  $|x_1|^2 = |x_2|^2 = 1/2$ . It then follows that  $(\langle A_3 \mathbf{x}, \mathbf{x} \rangle, \langle A_4 \mathbf{x}, \mathbf{x} \rangle) \neq (0, 0)$ . This proves that  $(1/2, 1/2, 0, 0) \notin W(\mathbf{A})$ . Hence,  $(0, 0, 0, 0) \in W_e(\mathbf{A}) \cap W(\mathbf{A})$  is not a star center of  $W(\mathbf{A})$  and  $W(\mathbf{A})$  is not convex.

To see that  $\text{cl}(W(\mathbf{A}))$  is convex, note that  $\mathbf{0} \in W_e(\mathbf{A})$ . Thus, by Theorem 3.1, for every  $\mathbf{b} \in \text{cl}(W(\mathbf{A}))$  we have  $t\mathbf{0} + (1-t)\mathbf{b} \in \text{cl}(W(\mathbf{A}))$  for any  $t \in [0, 1]$ .

Let  $\mathbf{B} = (B_1, B_2, B_3, B_4)$ , where  $B_1 = \text{diag}(0, 1, 0)$ ,  $B_2 = \text{diag}(0, 1, 0)$ ,

$$B_3 = [0] \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad B_4 = [0] \oplus \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix},$$

and  $\mathbf{C} = (C_1, C_2, C_3, C_4)$ , where  $C_1 = \text{diag}(1/3, 1/4, \dots) \oplus [0]$ ,  $C_2 = C_3 = C_4 = \text{diag}(0, 0, \dots) \oplus [0]$ . Then it is easy to verify that

$$W(\mathbf{B}) = \{(r, r, s, t) \in \mathbf{R}^4 : 4(r - 1/2)^2 + s^2 + t^2 \leq 1\}$$

and

$$W(\mathbf{C}) = \{(c, 0, 0, 0) : c \in [0, 1/3]\}$$

are both compact and convex. Hence,  $W(\mathbf{B} \oplus \mathbf{C}) = \text{conv}(W(\mathbf{B}) \cup W(\mathbf{C}))$  is compact and convex and

$$W(\mathbf{A}) \subseteq W(\mathbf{B} \oplus \mathbf{C}) \Rightarrow \text{cl}(W(\mathbf{A})) \subseteq W(\mathbf{B} \oplus \mathbf{C}).$$

On the other hand,  $\mathbf{B} \oplus \mathbf{C} = [0] \oplus \mathbf{A} \oplus [0]$ . Therefore,

$$W(\mathbf{B} \oplus \mathbf{C}) = \{t\mathbf{0} + (1-t)\mathbf{b} : \mathbf{b} \in W(\mathbf{A})\} \subseteq \text{cl}(W(\mathbf{A})).$$

So,  $\text{cl}(W(\mathbf{A})) = W(\mathbf{B} \oplus \mathbf{C})$  is convex.  $\square$

## 4 Other descriptions of $W_e(\mathbf{A})$

For  $\mathbf{c} = (c_1, \dots, c_m) \in \mathbf{R}^m$  and  $\mathbf{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ , let  $\mathbf{c} \cdot \mathbf{A} = \sum_{i=1}^m c_i A_i$ . Using the convexity of  $W_e(\mathbf{A})$ , we obtain additional equivalent formulations of  $W_e(\mathbf{A})$  in terms of  $\mathbf{c} \cdot \mathbf{A} \in \mathcal{S}(\mathcal{H})$  so that the joint behavior of  $A_1, \dots, A_m$  can be understood by their linear combinations. For  $A \in \mathcal{S}(\mathcal{H})$  and a positive integer  $k$ , let

$$\lambda_k(A) = \inf\{\max \sigma(A + F) : F \in \mathcal{S}(\mathcal{H}) \text{ with } \text{rank}(F) < k\}.$$

**Theorem 4.1** *Let  $\mathbf{A} \in \mathcal{S}(\mathcal{H})^m$  and  $\mathbf{a} = (a_1, \dots, a_m) \in \mathbf{R}^m$ . Then  $\mathbf{a} \in W_e(\mathbf{A})$  if and only if any one (and hence all) of the following conditions holds.*

- (1) *For every  $\mathbf{c} \in \mathbf{R}^m$ ,  $\mathbf{c} \cdot \mathbf{a} \in W_e(\mathbf{c} \cdot \mathbf{A})$ .*
- (2) *For every  $\mathbf{c} \in \mathbf{R}^m$ ,  $\mathbf{c} \cdot \mathbf{a} \in \cap\{\text{cl}(W(\mathbf{c} \cdot \mathbf{A} + F)) : F \in \mathcal{F}(\mathcal{H}) \cap \mathcal{S}(\mathcal{H})\}$ .*
- (3) *For every  $\mathbf{c} \in \mathbf{R}^m$ , there is an orthonormal sequence of vectors*

$$\{\mathbf{x}_n\}_{n=1}^\infty \in \mathcal{H} \quad \text{such that} \quad \lim_{n \rightarrow \infty} \langle \mathbf{c} \cdot \mathbf{A} \mathbf{x}_n, \mathbf{x}_n \rangle = \mathbf{c} \cdot \mathbf{a}.$$

- (4) *For every  $\mathbf{c} \in \mathbf{R}^m$ , there is a sequence of unit vectors  $\{\mathbf{x}_n\}_{n=1}^\infty \in \mathcal{H}$  such that  $\{\mathbf{x}_n\}_{n=1}^\infty$  converges weakly to  $\mathbf{0}$  in  $\mathcal{H}$  and*

$$\lim_{n \rightarrow \infty} \langle \mathbf{c} \cdot \mathbf{A} \mathbf{x}_n, \mathbf{x}_n \rangle = \mathbf{c} \cdot \mathbf{a}.$$

- (5) *For every  $\mathbf{c} \in \mathbf{R}^m$ , there is an infinite-dimensional projection  $P \in \mathcal{S}(\mathcal{H})$  such that  $P(\mathbf{c} \cdot \mathbf{A} - \mathbf{c} \cdot \mathbf{a}I)P \in \mathcal{K}(\mathcal{H})$ .*
- (6) *For every  $\mathbf{c} \in \mathbf{R}^m$  and  $k \geq 1$ ,  $\lambda_k(\mathbf{c} \cdot \mathbf{A} - \mathbf{c} \cdot \mathbf{a}I) \geq 0$ .*

*Proof.* By the convexity of  $W_e(\mathbf{A})$ , we can apply the separation theorem to Theorem 2.1 to show that  $\mathbf{a} \in W_e(\mathbf{A})$  if and only if any one of the conditions (1) to (5) holds.

To prove the equivalence of condition (6), suppose  $\mathbf{a} \in \mathbf{R}^m$ . Without loss of generality, we may assume that  $\mathbf{a} = \mathbf{0}$ . Suppose  $\mathbf{0}$  satisfies condition (6). Then for every  $\mathbf{c} \in \mathbf{R}^m$  and  $F \in \mathcal{F}(\mathcal{H}) \cap \mathcal{S}(\mathcal{H})$  with  $\text{rank } F = k$ , we have

$$\lambda_1(\mathbf{c} \cdot \mathbf{A} + F) \geq \lambda_{k+1}(\mathbf{c} \cdot \mathbf{A}) \geq 0 \quad \text{and} \quad \lambda_1(-(\mathbf{c} \cdot \mathbf{A} + F)) \geq \lambda_{k+1}(-\mathbf{c} \cdot \mathbf{A}) \geq 0.$$

Hence,  $\mathbf{c} \cdot \mathbf{0} = 0 \in \text{cl}(W(\mathbf{c} \cdot \mathbf{A} + F))$ . Therefore, condition (2) is satisfied.

Conversely, if  $\mathbf{0}$  does not satisfy condition (6), then there exist  $\mathbf{c} \in \mathbf{R}^m$  and  $k \geq 1$  such that  $\lambda_k(\mathbf{c} \cdot \mathbf{A}) < 0$ . Thus there exists  $F \in \mathcal{F}(\mathcal{H}) \cap \mathcal{S}(\mathcal{H})$  such that  $\mathbf{c} \cdot \mathbf{A} + F < 0$  and  $\mathbf{0}$  does not satisfy condition (2).  $\square$

Let  $\mathbf{A} \in \mathcal{S}(\mathcal{H})^m$ . Although the set  $\text{cl}(W(\mathbf{A}))$  may not be convex if  $m \geq 4$ , we have the following analog of the separation theorem for a convex set.

**Theorem 4.2** *Let  $\mathbf{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$  and  $\mathbf{d} = (d_1, \dots, d_m) \in \mathbf{R}^m$ . Then  $\mathbf{d} \notin W_e(\mathbf{A})$  if and only if any one (and hence all) of the following conditions holds..*

- (a) *There exists  $\mathbf{K} \in \mathcal{K}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m$  such that  $\mathbf{d} \notin \text{cl}(W(\mathbf{A} + \mathbf{K}))$ .*
- (b) *There exists  $\mathbf{F} \in \mathcal{F}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m$  such that  $\mathbf{d} \notin \text{conv}(\text{cl}(W(\mathbf{A} + \mathbf{F})))$ .*
- (c) *There exist  $F \in \mathcal{F}(\mathcal{H}) \cap \mathcal{S}(\mathcal{H})$ ,  $r > 0$  and  $\mathbf{c} = (c_1, \dots, c_m) \in \mathbf{R}^m$  such that*

$$\left( \sum_{i=1}^m c_i(A_i - d_i I) \right) + F > r I_{\mathcal{H}}. \quad (4)$$

*Proof.* For simplicity, replace  $(A_1, \dots, A_m)$  by  $(A_1 - d_1 I, \dots, A_m - d_m I)$  and assume that  $\mathbf{d} = (0, \dots, 0)$ .

(c) $\Rightarrow$ (b). Suppose (c) holds. We may perturb  $(c_1, \dots, c_m)$  so that  $c_j \neq 0$  for all  $j \in \{1, \dots, m\}$  so that condition (4) still holds true. In particular, we have  $c_1 \neq 0$ . Then let  $\mathbf{F} = (F/c_1, 0, \dots, 0)$ . We have  $\mathbf{c} \cdot \mathbf{a} > r > 0$  for all  $\mathbf{a} \in W(\mathbf{A} + \mathbf{F})$ . Therefore,  $\mathbf{0} \notin \text{conv}(\text{cl}(W(\mathbf{A} + \mathbf{F})))$ .

Clearly, we have (b)  $\Rightarrow$  (a), which implies that  $\mathbf{0} \notin W_e(\mathbf{A})$ .

Finally, suppose  $\mathbf{0} \notin W_e(\mathbf{A})$ . Then by Theorem 4.1 (2), there exist a real vector  $\mathbf{c} = (c_1, \dots, c_m)$  and  $F \in \mathcal{F}(\mathcal{H}) \cap \mathcal{S}(\mathcal{H})$  such that  $0 = \mathbf{c} \cdot \mathbf{0} \notin \text{cl}(W(\mathbf{c} \cdot \mathbf{A} + F))$ . Since  $\text{cl}(W(\mathbf{c} \cdot \mathbf{A} + F))$  is a closed interval  $[s, t]$  of  $\mathbf{R}$ , we may assume that  $0 < s \leq t$ . Let  $r = s/2$ , we have  $(\sum_{i=1}^m c_i A_i) + F > r I_{\mathcal{H}}$ . Hence, (c) holds.  $\square$

Let  $\Omega = \{\mathbf{c} \in \mathbf{R}^m : \langle \mathbf{c}, \mathbf{c} \rangle = 1\}$ . By Theorem 4.2, we have the following result showing that  $W_e(\mathbf{A})$  can be expressed as the intersection of half spaces.

**Corollary 4.3** *Let  $\mathbf{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ . Then*

$$\begin{aligned} W_e(\mathbf{A}) &= \bigcap_{\mathbf{c} \in \Omega} \{\mathbf{d} \in \mathbf{R}^m : \langle \mathbf{c}, \mathbf{d} \rangle \leq \max W_e(\mathbf{c} \cdot \mathbf{A})\} \\ &= \{\mathbf{d} \in \mathbf{R}^m : \langle \mathbf{c}, \mathbf{d} \rangle \in W_e(\mathbf{c} \cdot \mathbf{A}) \text{ for all } \mathbf{c} \in \Omega\}. \end{aligned}$$

For  $A \in \mathcal{B}(\mathcal{H})$ , let  $\sigma_e(A) = \cap\{\sigma(A + K) : K \in \mathcal{K}(\mathcal{H})\}$  bet the essential spectrum of  $A$ . Then for  $A \in \mathcal{S}(\mathcal{H})$ , we have

$$W_e(A) = \mathbf{conv}\sigma_e(A).$$

Thus, one may replace  $\max W_e(\mathbf{c} \cdot \mathbf{A})$  by  $\max \sigma_e(\mathbf{c} \cdot \mathbf{A})$  in Corollary 4.3.

**Corollary 4.4** *Let  $\mathbf{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ . If  $\mathbf{d} \notin \mathbf{cl}(W(\mathbf{A}))$ , then for any  $i \in \{1, \dots, m\}$  there exists  $F \in \mathcal{F}(\mathcal{H}) \cap \mathcal{S}(\mathcal{H})$  such that  $\mathbf{d} \notin \mathbf{conv}(\mathbf{cl}(W(\tilde{\mathbf{A}})))$ , where  $\tilde{\mathbf{A}} = (A_1, \dots, A_{i-1}, A_i + F, A_{i+1}, \dots, A_m)$ .*

*Proof.* If  $\mathbf{d} \notin \mathbf{cl}(W(\mathbf{A}))$ , then  $\mathbf{d} \notin W_e(\mathbf{A})$ . The result readily follows from the arguments in the last paragraph in proof of Theorem 4.2.  $\square$

It follows from Theorem 2.1 that the intersection of the non-convex sets  $\mathbf{cl}(W(\mathbf{A} + \mathbf{K}))$ , which equals  $W_e(\mathbf{A})$ , is a convex set. By Theorem 4.2 and Corollary 4.4, we see that one can replace  $\mathbf{cl}(W(\mathbf{A} + \mathbf{K}))$  by its convex hull for the intersection to obtain the same convex set  $W_e(\mathbf{A})$ . It is known that for any  $\mathbf{B} = (B_1, \dots, B_m) \in \mathcal{B}(\mathcal{H})^m$ ,

$$\mathbf{conv}(\mathbf{cl}(W(\mathbf{B}))) = \{(f(B_1), \dots, f(B_m)) : f \in \Omega\},$$

where  $\Omega$  is the set of linear functionals  $f$  on  $\mathcal{B}(\mathcal{H})$  satisfying  $1 = f(I) = \max\{f(X) : X \in \mathcal{B}(\mathcal{H}), \|X\| \leq 1\}$ ; for example, see [10, 11]. So, it is easier to determine  $\mathbf{conv}(\mathbf{cl}(W(\mathbf{A} + \mathbf{K})))$  than  $\mathbf{cl}(W(\mathbf{A} + \mathbf{K}))$ . In fact, we have the following.

**Corollary 4.5** *Let  $\mathbf{A} \in \mathcal{S}(\mathcal{H})^m$  and  $i \in \{1, \dots, m\}$ . Then*

$$\begin{aligned} & W_e(\mathbf{A}) \\ &= \cap\{\mathbf{cl}(W(\mathbf{A} + \mathbf{F})) : \mathbf{F} \in \{0\}^{i-1} \times (\mathcal{F}(\mathcal{H}) \cap \mathcal{S}(\mathcal{H})) \times \{0\}^{m-i}\} \\ &= \cap\{\mathbf{conv}(\mathbf{cl}(W(\mathbf{A} + \mathbf{F}))) : \mathbf{F} \in \{0\}^{i-1} \times (\mathcal{F}(\mathcal{H}) \cap \mathcal{S}(\mathcal{H})) \times \{0\}^{m-i}\}. \end{aligned}$$

*Proof.* Let  $\mathbf{F} \in \{0\}^{i-1} \times (\mathcal{F}(\mathcal{H}) \cap \mathcal{S}(\mathcal{H})) \times \{0\}^{m-i}$ . Clearly, we have

$$W_e(\mathbf{A}) \subseteq \mathbf{cl}(W(\mathbf{A} + \mathbf{F})) \subseteq \mathbf{conv}(\mathbf{cl}(W(\mathbf{A} + \mathbf{F}))).$$

So, we may take the intersection of the second and third sets over all  $\mathbf{F} \in \{0\}^{i-1} \times (\mathcal{F}(\mathcal{H}) \cap \mathcal{S}(\mathcal{H})) \times \{0\}^{m-i}$ , and get a set inclusion relation involving the three sets in the corollary. Finally, if  $\mathbf{d} \notin W_e(\mathbf{A})$ , then  $\mathbf{d}$  will not belong to the third set by Corollary 4.4. So, the third set is a subset of  $W_e(\mathbf{A})$ . Hence, the three sets in the corollary are equal.  $\square$

## 5 Additional Results

The following result shows that  $W_e(\mathbf{A})$  is unchanged under certain operations on  $\mathbf{A}$ .

**Theorem 5.1** *Let  $\mathbf{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ .*

- (a) *Suppose  $\mathcal{H}_1$  is a closed subspace of  $\mathcal{H}$  such that  $\mathcal{H}_1^\perp$  is finite dimensional. If  $X : \mathcal{H}_1 \rightarrow \mathcal{H}$  is such that  $X^*X = I_{\mathcal{H}_1}$ , then*

$$W_e(\mathbf{A}) = W_e(X^*A_1X, \dots, X^*A_mX).$$

- (b) *For each  $j \in \{1, \dots, m\}$ , suppose  $P_j : \mathcal{H} \rightarrow \mathcal{H}$  is an orthogonal projection such that  $I - P_j$  has finite rank. Then*

$$W_e(\mathbf{A}) = W_e(P_1A_1P_1, \dots, P_mA_mP_m).$$

*Proof.* Using the formulation of  $W_e(\mathbf{A})$  in Theorem 2.1, one readily shows that the set equalities in (a) and (b) hold.  $\square$

We will establish some additional relationships between the sets  $W_e(\mathbf{A})$  and  $W(\mathbf{A})$ . The next theorem generalizes the results in [29] and [14].

**Theorem 5.2** *Let  $\mathbf{A} \in \mathcal{S}(\mathcal{H})^m$ . Then  $W_e(\mathbf{A}) = \mathbf{cl}(W(\mathbf{A}))$  if and only if  $\text{Ext}(W(\mathbf{A})) \subseteq W_e(\mathbf{A})$ .*

*Proof.* If  $W_e(\mathbf{A}) = \mathbf{cl}(W(\mathbf{A}))$ , then we have

$$\text{Ext}(W(\mathbf{A})) \subseteq W(\mathbf{A}) \subseteq W_e(\mathbf{A}).$$

Conversely, if  $\text{Ext}(W(\mathbf{A})) \subseteq W_e(\mathbf{A})$ , then by (P6), we have

$$\text{Ext}(\mathbf{cl}(W(\mathbf{A}))) \subseteq W_e(\mathbf{A}).$$

Hence,

$$\mathbf{cl}(W(\mathbf{A})) \subseteq \mathbf{conv}(\text{Ext}(\mathbf{cl}(W(\mathbf{A})))) \subseteq \mathbf{conv}(W_e(\mathbf{A})) = W_e(\mathbf{A}).$$

Since  $W_e(\mathbf{A}) \subseteq \mathbf{cl}(W(\mathbf{A}))$ , we have  $W_e(\mathbf{A}) = \mathbf{cl}(W(\mathbf{A}))$ .  $\square$

For  $k \geq 1$ , let  $I_k$  denotes the  $k \times k$  identity matrix. Then for  $\mathbf{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ , we have  $\mathbf{A} \otimes I_k = (A_1 \otimes I_k, \dots, A_m \otimes I_k) \in \mathcal{S}(\underbrace{\mathcal{H} \oplus \dots \oplus \mathcal{H}}_{k\text{-copies}})^m$ .

Similarly, let  $I_\infty$  denotes the identity operator acting on  $\ell_2$ . Then for  $\mathbf{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ , we have  $\mathbf{A} \otimes I_\infty = (A_1 \otimes I_\infty, \dots, A_m \otimes I_\infty) \in \mathcal{S}(\underbrace{\mathcal{H} \oplus \mathcal{H} \oplus \dots}_{\text{infinite copies}})^m$ .

**Theorem 5.3** Let  $\mathbf{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ . Then for any positive integer  $k > \sqrt{m} - 1$ ,

$$W(\mathbf{A} \otimes I_k) = \mathbf{conv}(W(\mathbf{A})).$$

Moreover, we have

$$W_e(\mathbf{A} \otimes I_\infty) = \mathbf{cl}(\mathbf{conv}(W(\mathbf{A}))).$$

*Proof.* Suppose  $k > \sqrt{m} - 1$ . By the result in [34], every  $\mathbf{a} \in \mathbf{conv}(W(\mathbf{A}))$  can be written as  $\mathbf{a} = \sum_{j=1}^k t_j \langle \mathbf{A} \mathbf{x}_j, \mathbf{x}_j \rangle$  for some unit vectors  $\mathbf{x}_1, \dots, \mathbf{x}_k \in \mathcal{H}$ . Thus, for  $\mathbf{x} = (\sqrt{t_1} \mathbf{x}_1, \dots, \sqrt{t_k} \mathbf{x}_k) \in \mathcal{H} \oplus \dots \oplus \mathcal{H}$ , we have  $\langle \mathbf{A} \otimes I_k \mathbf{x}, \mathbf{x} \rangle = \mathbf{a}$ . Conversely, if  $\mathbf{a} = \langle \mathbf{A} \otimes I_k \mathbf{x}, \mathbf{x} \rangle \in W(\mathbf{A} \otimes I_k)$ , one can decompose the unit vector  $\mathbf{x}$  into  $k$  parts  $\mathbf{y}_1, \dots, \mathbf{y}_k$  according to the structure of  $\mathcal{H} \otimes I_k$ . Then

$$\mathbf{a} = \sum_{j=1}^k \|\mathbf{y}_j\|^2 \langle A \mathbf{y}_j / \|\mathbf{y}_j\|, \mathbf{y}_j / \|\mathbf{y}_j\| \rangle \in \mathbf{conv}(W(\mathbf{A})).$$

If  $\mathbf{a} \in \mathbf{cl}(\mathbf{conv}(W(\mathbf{A})))$ , then there is a sequence of unit vectors  $\{\mathbf{x}_n\}$  in  $\mathcal{H}$  such that  $\langle \mathbf{A} \mathbf{x}_n, \mathbf{x}_n \rangle \rightarrow \mathbf{a}$ . Let

$$\tilde{\mathbf{x}}_n = (\underbrace{0, \dots, 0}_{n-1 \text{ terms}}, \mathbf{x}_n, 0, \dots) \in \mathcal{H} \oplus \mathcal{H} \oplus \mathcal{H} \oplus \dots.$$

Then  $\{\tilde{\mathbf{x}}_n\}$  is an orthonormal sequence in  $\mathcal{H} \oplus \mathcal{H} \oplus \mathcal{H} \oplus \dots$  such that  $\langle \mathbf{A} \otimes I_\infty \tilde{\mathbf{x}}_n, \tilde{\mathbf{x}}_n \rangle \rightarrow \mathbf{a}$ . Therefore,  $\mathbf{a} \in W_e(\mathbf{A} \otimes I_\infty)$ . Since

$$\begin{aligned} W_e(\mathbf{A} \otimes I_\infty) &\subseteq \mathbf{cl}(W(\mathbf{A} \otimes I_\infty)) \\ &= \mathbf{cl}\left(\bigcup_{k=1}^{\infty} W(\mathbf{A} \otimes I_k)\right) \subseteq \mathbf{cl}(\mathbf{conv}(W(\mathbf{A}))), \end{aligned}$$

we get the reverse inclusion.  $\square$

**Corollary 5.4** Let  $S$  be a compact convex subset of  $\mathbf{R}^m$ . Then there are  $\mathbf{A}, \tilde{\mathbf{A}} \in \mathcal{S}(\mathcal{H})^m$  with  $\mathcal{H} = \ell^2$  such that  $W(\mathbf{A})$  is convex and

$$W(\mathbf{A}) \subseteq S = \mathbf{cl}(W(\mathbf{A})) = W_e(\tilde{\mathbf{A}}).$$

*Proof.* For  $j = 1, \dots, m$ , let  $A_j = \text{diag}(a_{1j}, a_{2j}, \dots)$  act on  $\ell^2$  with the standard canonical basis  $\{e_n : n \geq 1\}$  such that  $\{(a_{i1}, a_{i2}, \dots, a_{im}) : i \geq 1\}$  is a dense subset of  $S$ . Then for  $\mathbf{A} = (A_1, \dots, A_m)$ , we have

$$W(\mathbf{A}) = \mathbf{conv}\{(a_{i1}, a_{i2}, \dots, a_{im}) : i \geq 1\}$$

is convex, and  $\tilde{\mathbf{A}} = \mathbf{A} \otimes I_\infty$  will satisfy the assertion by Theorem 5.3.  $\square$

## References

- [1] Y.S. Abramov, On the geometric properties of the joint spectrum of a family of self-adjoint operators, *Studia Math.* 61 (1977), 55-62.
- [2] J.A. Anderson, On compact perturbations of operators, *Canad. J. Math.* 26 (1974), 247–250.
- [3] S.C. Arora and R. Kumar, Joint essential numerical range, *Glas. Mat. Ser. III* 18(38) (1983), 317–320.
- [4] Y.H. Au-Yeung and Y.T. Poon, A Remark on the Convexity and Positive Definiteness Concerning Hermitian Matrices, *SEA Bull. Math.* 3 (1979), 85-92.
- [5] Y.H. Au-Yeung and N.K. Tsing, An extension of the Hausdorff-Toeplitz theorem on the numerical range, *Proc. Amer. Math. Soc.* 89 (1983), 215–218.
- [6] M. Barraa, Essential numerical range of elementary operators, *Proc. Amer. Math. Soc.* 133 (2005), 1723–1726
- [7] M. Barraa and V. Müller, On the essential numerical range, *Acta Sci. Math. (Szeged)* 71 (2005), no. 1-2, 285–298.
- [8] S.K. Berberian and G.H. Orland, On the closure of the numerical range of an operator, *Proc. Amer. Math. Soc.* 18 (1967), 499-503.
- [9] P. Binding and C.K. Li, Joint ranges of Hermitian matrices and simultaneous diagonalization, *Linear Algebra Appl.* 151 (1991), 157-168.
- [10] F.F. Bonsall and J.Duncan, *Numerical Ranges of Operators on Normed Spaces and Elements of Normed Algebras*, London Math. Soc. Lecture Notes Series 2, Cambridge University Press, Cambridge, 1971.
- [11] F.F. Bonsall and J. Duncan, *Numerical ranges, II*, Cambridge University Press, Cambridge, 1973.
- [12] J.J. Buoni and B.L. Wadhwa, On joint numerical ranges, *Pacific J. Math* 77 (1978), 303-306.

- [13] G.F. Cao, On the joint essential spectrum, the joint essential numerical range and the boundary of joint numerical range, *J. Math. Res. Exposition* 10 (1990), 25–31.
- [14] M. Cho and M. Takagunchi, Boundary points of joint numerical ranges, *Pacific J. Math.* 95 (1981), 27–35.
- [15] A.T. Dash, Joint numerical range, *Glasnik Mat.* 7 (1972), 75–81.
- [16] A.T. Dash, Joint spectra, *Studia Math* 45 (1973), 225–237.
- [17] A.T. Dash, Joint spectrum in the Calkin algebra, *Bulletin Amer. Math. Soc.* 81 (1975), 1083–1085.
- [18] T.M., El-Adawy, M.Z. Abd-Alla and H.A. Atia, On the joint essential and joint maximal essential centre valued range, *Proc. Math. Phys. Soc. Egypt No. 77* (2002), 29–49 (2003).
- [19] M.K.H. Fan and A.L. Tits, On the generalized numerical range, *Linear and Multilinear Algebra* 21 (1987), 313–320.
- [20] P. Fan and C.K. Fong and D.A. Herrero, On zero-diagonal operators and traces, *Proc. Amer. Math. Soc.* 99 (1987), 445–451.
- [21] P.A. Fillmore, J.G. Stampfli and J.P. Williams, On the essential numerical range, the essential spectrum, and a problem of Halmos, *Acta Sci. Math. (Szeged)* 33 (1972), 179–192
- [22] K.E. Gustafson and D.K.M. Rao, *Numerical ranges: The field of values of linear operators and matrices*, Springer, New York, 1997.
- [23] E. Gutkin, E.A. Jonckheere and M. Karow, Convexity of the joint numerical range: topological and differential geometric viewpoints, *Linear Algebra Appl.* 376 (2004), 143–171.
- [24] P.R. Halmos, *A Hilbert Space Problem Book*, Second edition, Graduate Texts in Mathematics, 19, Encyclopedia of Mathematics and its Applications, Springer-Verlag, New York, 1982.
- [25] R.A. Horn and C.R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, 1991.
- [26] G. Ji, N. Liu, and Z. Li, Essential numerical range and maximal numerical range of the Aluthge transform, *Linear and Multilinear Algebra* 55 (2007), 315–322.

- [27] M.S. Jones and H.P. Rogosinski, Inclusion relations involving the  $k$ -numerical range and the essential numerical range. Special Issue: The numerical range and numerical radius, *Linear and Multilinear Algebra* 37 (1994), 161–173.
- [28] S.P. Juneja, On extreme points of the joint numerical range of commuting normal operators, *Pacific J. Math.* 67 (1976), 473-476.
- [29] J.S. Lancaster, The boundary of the numerical range, *Proc. Amer. Math. Soc.* 49 (1975), 393-398.
- [30] D.A. Legg and D.W. Townsend, Essential numerical range in  $B(l_1)$ , *Proc. Amer. Math. Soc.* 81 (1981), 541–545.
- [31] C.K. Li and Y.T. Poon, Convexity of the joint numerical range, *SIAM J. Matrix Analysis Appl.* 21 (1999), 668-678.
- [32] B. Magajna, On the essential numerical range of a generalized derivation, *Proc. Amer. Math. Soc.* 99 (1987), 86–92.
- [33] Y. Ogata, Local distinguishability of quantum states in infinite-dimensional systems, *J. Phys. A* 39 (2006), no. 12, 3059–3069.
- [34] Y.T. Poon, On the convex hull of the multiform numerical range, *Linear and Multilinear Algebra* 37 (1994), 221-223.
- [35] Y.T. Poon, Generalized numerical ranges, joint positive definiteness and multiple eigenvalues, *Proc. Amer. Math. Soc.* 125 (1997), no. 6, 1625–1634.
- [36] M. Takagunchi and M. Cho, The joint numerical range and the joint essential numerical range, *Sci. Prp. Hirosaki Univ.* 27 (1980), 6-8.
- [37] J.P. Williams, The numerical range and the essential numerical range, *Proc. Amer. Math. Soc.* 66 (1977), 185-186.