

# MATH 451/551

## Chapter 5. Common Continuous Distribution

### 5.5 Normal Distribution

GuanNan Wang  
[gwang01@wm.edu](mailto:gwang01@wm.edu)



# Motivating Examples



## Motivating Examples

- ▶ Many random variables that arise in practice have bell-shaped distributions. Examples include
  - ▶ cholesterol levels of 50-year-old men
  - ▶ heights of adult women
  - ▶ weights of newborn babies
  - ▶ crop yields
  - ▶ ball bearing diameters

# Normal Distribution



## Normal Distribution

A continuous random variable  $X$  with probability density function

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad -\infty < x < \infty$$

for some real constants  $\mu$  and  $\sigma > 0$  is a  $N(\mu, \sigma^2)$  random variable.



# Mean



# Variance



# R Functions



## R Functions

Function	Returned Value
<code>dnorm(x, <math>\mu</math>, <math>\sigma</math>)</code>	calculates the probability density function $f(x)$
<code>pnorm(x, <math>\mu</math>, <math>\sigma</math>)</code>	calculates the cumulative distribution function $F(x)$
<code>qnorm(u, <math>\mu</math>, <math>\sigma</math>)</code>	calculates the percentile (quantile) $F^{-1}(u)$
<code>rnorm(m, <math>\mu</math>, <math>\sigma</math>)</code>	generates $m$ random variates


# Properties



- Shorthand:  $X \sim N(\mu, \sigma^2)$
- Since  $f(x)$  symmetric, mean, median, and mode all equal  $\mu$
- The pdf has inflection points at  $\mu \pm \sigma$
- Special case: standard normal distribution  $Z \sim N(0, 1)$
- The cumulative distribution function of  $X \sim N(\mu, \sigma^2)$  is

$$\int_{-\infty}^x \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(w-\mu)^2}{2\sigma^2}} dw, \quad -\infty < x < \infty.$$

- Some authors use  $\Phi(x)$ , rather than  $F(x)$ , for the standard normal cdf
- Empirical Rule (68–95–99.7 Rule)



# Theorem 5.3



## Theorem 5.3

If  $X \sim N(\mu, \sigma^2)$  then  $Y = aX + b \sim N(a\mu + b, a^2\sigma^2)$  for real-valued constants  $a \neq 0$  and  $b$ .

# Theorem 5.4



## Theorem 5.4

If  $X \sim N(\mu, \sigma^2)$  then  $Z = \frac{X-\mu}{\sigma} \sim N(0, 1)$ .

# Theorem 5.5



10

## Theorem 5.5

If  $X \sim N(\mu, \sigma^2)$  then  $Y = \left(\frac{X-\mu}{\sigma}\right)^2 \sim \chi^2(1)$ .

# Example 1



## Example 1

IQ scores are normally distributed with population mean 100 and population standard deviation 15.

1. Find the probability that an IQ score is less than 130.
2. Find the probability that an IQ score falls between 80 and 110.
3. Find the 99th percentile of the distribution of IQ scores.

## Example 2



## Example 2

If  $X \sim N(\mu, \sigma^2)$ , find  $E(|X - \mu|)$ .

# Thank You



# THANK YOU!

